



Acelepryn[®]

Lawn Grub and Army Worm Control

syngenta[®]

ACTIVE CONSTITUENT: 200 g/L CHLORANTRANILIPROLE

For the control of turf pests including Lawn Army Worm and Lawn Curl Grubs. Suitable for all lawn varieties

Syngenta Australia Pty Ltd

Level 1, 2 Lyonpark Road, Macquarie Park NSW 2113

***In a transport emergency only, dial 000, Police or Fire Brigade
For specialist advice in an emergency only, call 1800 033 111
(24 hours)***

APVMA Approval No.: 91647/132981



HOW TO USE

ACELEPRYN® Lawn Grub and Army Worm Control delivers excellent preventative control against a large range of common lawn pests. Optimal results are achieved when applied early in the season at the appearance of overwintering adult pests. The long residual performance of ACELEPRYN® Lawn Grub and Army Worm Control will provide up to 6 months protection at the higher application rates. Curative control of pest outbreaks later in the season can also be achieved using the higher recommended application rates.

Restraints

DO NOT apply when the soil is saturated with water as adequate distribution of the product into the soil will not be achieved.

Situation	Pest	Rate	Comments
All Lawn Varieties	Lawn Curl Grub (Beetle larvae including: African Black Beetle larvae Argentinian Scarab larvae)	Garden Sprayer Use 8 – 15 mL per 5-10 L of water per 100 m ² Watering Can Mix 1.5 mL in 9 L of water and evenly apply to 10 m ²	African Black Beetle larvae Apply before peak egg hatch for maximum control (typically mid-September). Argentinian Scarab larvae Apply before or at peak egg hatch for maximum control (typically mid-December). Apply the higher rate for: <ul style="list-style-type: none"> • up to 6 months residual protection • cases of high pest pressure • late in season when mature grubs are present Lightly water in immediately after application when applying by garden sprayer.
	Argentine Stem Weevil larvae Billbug larvae		African Black Beetle larvae Apply before peak egg hatch for maximum control (typically mid-September). Argentinian Scarab larvae Apply before or at peak egg hatch for maximum control (typically mid-December). Apply the higher rate for: <ul style="list-style-type: none"> • up to 6 months residual protection • cases of high pest pressure • late in season when mature grubs are present Lightly water in immediately after application when applying by garden sprayer.
	Caterpillars including Lawn Armyworm, Black Cutworm, and Sod Webworm	Garden Sprayer Use 8 mL per 5-10 L of water per 100 m ²	To ensure optimum control, delay watering or mowing for 24 hours after application.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED

WITHHOLDING PERIODS

DO NOT GRAZE TREATED TURF OR LAWN; OR FEED TURF OR LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK

HOW TO PREPARE

Mix ACELEPRYN® Lawn Grub and Army Worm Control concentrate as per the instructions below into a diluted mix ready for application. If dilute solution is left standing, make sure you shake/agitate before use.

Mixing – Garden Sprayer

1. Prior to opening, shake ACELEPRYN® bottle vigorously.
2. Add half the required amount of water to your garden sprayer.
3. Add the required quantity of ACELEPRYN® to the half full garden sprayer.
4. Add the remaining water required and ensure closed before agitating sprayer.
5. Only mix sufficient chemical for application area and for the day of mixing.

Mixing – Watering Can

1. Prior to opening, shake ACELEPRYN® bottle vigorously.
2. Add a quarter of the required amount of water to your watering can.
3. Add the required quantity of ACELEPRYN® to the watering can.
4. Add the remaining water required and use the process to agitate the mixture.
5. Only mix sufficient chemical for application area and for the day of mixing.

HOW TO APPLY

Measure lawn area before application to determine size and make up enough mixture to apply over area. Evenly spray or pour the mixed solution over the area to be treated. For optimal control of beetle larvae (Black beetle, Argentine scarab, Argentine stem weevil and Billbug) apply preventatively and lightly water in immediately after application, when applying by garden sprayer. When treating for caterpillars apply when pests or damage is observed. Irrigation and mowing should be delayed until 24 hours after application when treating for caterpillars. For more information regarding calibrating knapsack sprayers visit syngentaturf.com.au.

Thoroughly clean all equipment after application. Empty and thoroughly rinse equipment such as sprayer, hoses, nozzles and watering can with clean water.

CAUTION

Very toxic to aquatic life. DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds. No re-entry period is required. Re-entry is permitted immediately after spraying.

STORAGE AND DISPOSAL:

KEEP OUT OF REACH OF CHILDREN

Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

This product must be used strictly as directed. Not to be used for any use or in any manner contrary to the label.

Batch No

BARCODE

Date of Manufacture